AD					

Award Number: W81XWH-04-1-0084

TITLE: Identification of a Protein for Prostate-Specific Infection

PRINCIPAL INVESTIGATOR: Shen Pang

CONTRACTING ORGANIZATION: University of California
Los Angeles, CA 90095

REPORT DATE: December 2006

TYPE OF REPORT: Annual Summary

PREPARED FOR: U.S. Army Medical Research and Materiel Command

Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

		<u>.</u>
REPORT DO	CUMENTATION PAGE	Form Approved OMB No. 0704-0188
Public reporting burden for this collection of information is data needed, and completing and reviewing this collection this burden to Department of Defense, Washington Headqu	estimated to average 1 hour per response, including the time for reviewin of information. Send comments regarding this burden estimate or any of uarters Services, Directorate for Information Operations and Reports (07, any other provision of law, no person shall be subject to any penalty for	ing instructions, searching existing data sources, gathering and maintaining the ther aspect of this collection of information, including suggestions for reducing 04-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-failing to comply with a collection of information if it does not display a currently
1. REPORT DATE (DD-MM-YYYY)	2. REPORT TYPE	3. DATES COVERED (From - To)
01/12/06	Annual Summary	1 Dec 2005 - 30 Nov 2006
4. TITLE AND SUBTITLE Identification of a Protein for I	5a. CONTRACT NUMBER	
		5b. GRANT NUMBER
		W81XWH-04-1-0084
		5c. PROGRAM ELEMENT NUMBER
6. AUTHOR(S) Shen Pang	5d. PROJECT NUMBER	
		5e. TASK NUMBER
		5f. WORK UNIT NUMBER
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)	8. PERFORMING ORGANIZATION REPORT NUMBER
University of California		
Los Angeles, CA 90095		
9. SPONSORING / MONITORING AGENCY U.S. Army Medical Research		10. SPONSOR/MONITOR'S ACRONYM(S)
Fort Detrick, Maryland 21	702-5012	
		11. SPONSOR/MONITOR'S REPORT NUMBER(S)
12. DISTRIBUTION / AVAILABILITY STAT	EMENT	
	se; Distribution Unlimited	
13. SUPPLEMENTARY NOTES Original contains colore	d plates: ALL DTIC reproductions	will be in black and white.
vector. We expect that using to cytotoxic genes into prostate concer cells in patients. During lentiviral vector envelop gene a vector that have peptide YDSV cells in 24-well plates. The concused to infect LNCaP cells as the	his protein to modify various gene thancer cells using systemic treatment, at the third year, we inserted the sequence and used one of the modified membran LALSAALQATR (P2) modified gp41 trol vector that does not have gp41-P2	to generate infection-specific gene therapy herapy vectors, we can specifically deliver and eventually eradicate metastatic prostate es that encode these two peptides to modify e proteins to target prostate cells. The viral envelope protein was used to infect LNCaP envelope protein on viral surface was also at with the gp41-P2 envelope protein on the to 70%.
NOT PROVIDED		

17. LIMITATION

OF ABSTRACT

18. NUMBER

6

OF PAGES

16. SECURITY CLASSIFICATION OF:

b. ABSTRACT

c. THIS PAGE

a. REPORT

Annual

19a. NAME OF RESPONSIBLE PERSON

19b. TELEPHONE NUMBER (include area

Shen Pang

(310) 825-0863

code)

Table of Contents

Cover	1
SF 298	2
Table of Contents	3
Introduction	4
Body	4
Key Research Accomplishments	6
Reportable Outcomes	
Conclusions€	;
References	
Appendices	

Title: Identification of a Protein for Prostate-Specific Infection

PI: Shen Pang

Start date: Dec. 1, 2003

Grant number: W81XWH-04-1-0084

Introduction:

Using gene therapy vectors to eradicate prostate cells is extensively studied. It is expected that if we can use a target-specific gene therapy to deliver cytotoxic genes into metastatic prostate cancer cells in patients, we can eradicate these malignant cells and cure the patients. To find an approach to generate infection-specific vectors is critical for target-specific cancer cell eradication. Because viral vectors infect cells by the binding of their surface proteins to their receptors on the target cells, it is expected that infection-specific vectors can be generated by modifying their surface proteins with proteins that can specifically bind to target cells. In our Preliminary Studies, we found that the lentiviral vector (an HIV-based retroviral vector) generated from a cell line derived from human oral tissue (HOT) can specifically infect LNCaP prostate cancer cells. Virus generated from other cell lines has much lower infectivity to LNCaP cells (200 to 1200-fold lower). These results strongly suggest that the HOT cells express a very specific membrane protein that can be picked by lentiviral vectors to modify their envelope. With such protein on its envelope, the viral vectors are able to specifically infect LNCaP cells.

If this is the case, we expect that a protein for generating infection-specific vector can be obtained by screen the cDNA library of the HOT cell line.

Body:

The approved SOW is listed below.

- Task 1. To prepare cDNA library in eukaryotic gene expression vectors (months 1-8).
- Task 2. To perform first-round screening to identify the cDNA clone groups that contain the cDNA clones encoding the protein responsible for tissue-specific infection. (months 9-20)
- Task 3. To perform the second round of screening to identify the individual clones that contain the cDNA encoding the protein responsible for tissue-specific infection (months 21-24)
- Task 4. To sequence the identified gene (months 25-26)
- Task 5. To use sequence analysis to characterize the identified gene (months 27-28)
- Task 6. To generate deletions of the identified gene and to use them to confirm the functional domains of the identified gene (months 31-36)

To follow the schedule, we should complete all the Tasks. We have identified two peptides that can significantly increase tissue-specific gene delivery as we reported in the annual report of last year (Task 1-4). In the past year, we have inserted the sequences into a lentiviral vector to increase gene delivery efficiency.

Progress of our research:

During the third year, we have focused how to use the two peptides to modify lentiviral vectors, so that the generated lentiviral vectors are able to specifically target prostate cancer cells (Task 6).

We make progress following our SOW.

1. Insert sequences into lentiviral envelope protein gp41

As described in the annual report of last year, we have identified two peptides that can increase lentiviral infection of LNCaP prostate cells 2 to 4-fold. We hypothesize that by adding these two peptides to lentiviral envelope protein, we can make lentiviral vector to specifically infect LNCaP cells. As Figure 1 shown, gp41 protein is an envelope protein located on the surface of lentiviral vector. The N-terminus of this protein is extracellular. So it is possible to add our sequence to the N-terminus so that the added peptide sequences can interact with LNCaP surface protein. When the lentiviral vector attaches to the target cells, the other part of gp41 can induce the viral envelope to fuse with cell membrane and render the viral genome to enter the target cells.




Fig. 1. Strategy of our approach to use two peptides with sequence of RGCICRCIGRGCICRCIG or YDSVLALSAALQAT R to modify lentiviral envelope protein gp41. A) A diagram of a lentiviral vector. B) Attachment of our peptide to gp41 at the N-terminus. C) The structure of a plasmid that carries the gp41 fusion fusion peptide.

2. Using modified lentiviral vectors to infect LNCaP cells

We used co-transfection method to generate lentiviral vectors that carry the enhanced green fluorescent protein (EGFP). The viral vector that have peptide

YDSVLALSAALQATR (P2) modified gp41 envelope protein was used to infect LNCaP cells in 24-well plates. The control vector that does not have gp41-P2 envelope protein on viral surface was also used to infect LNCaP cells as the control. Our results demonstrated that with the gp41-P2 envelope protein on the surface of the viral vector, lentiviral vectors increase infectivity by 30% to 70% as shown in Fig. 2.




Fig. 2. Increased infectivity by peptide YDSVLALSAALQATR. LNCaP cells were infected by a EGFP-expression lentiviral vector with gp41-P2 envelope protein on viral surface or with no gp41-P2 (control) on viral surface. The left penal, Cells in 24-well plates were infected by viral vector of 100 ng p24 reading. The right penal, LNCaP cells infected by 50 and 150 ng of lentiviral

Key Research accomplishments:

We have fused one of our peptides into a lentiviral envelope protein and found that the addition of this peptide increased gene delivery efficiency

Conclusions:

We have inserted two peptide sequences into a lentiviral surface protein gp41 and tested one. Our results demonstrated that with such sequence, gene delivery efficiency can be enhanced. However, the increase is not very high. It is expected by using different way to modify the lentiviral envelope gene with these two peptides, we can increase gene delivery efficient even further. We will insert multiple copies of our peptide sequences into gp41 and to insert these peptide sequences into other gp41 locations. We expect that by these approaches, we can further increase gene delivery efficiency.